

A CHARACTERIZATION OF THE ROGERS Q-HERMITE POLYNOMIALS

WALEED A. AL-SALAM

ABSTRACT. In this paper we characterize the Rogers q-Hermite polynomials as the only orthogonal polynomial set which is also $\mathcal{D}_q$ -Appell where $\mathcal{D}_q$ is the Askey-Wilson finite difference operator.

1. INTRODUCTION

Appell polynomials sets $\{P_n(x)\}$ are generated by the relation

$$(1.1) \quad A(t)e^{xt} = \sum_{n=0}^{\infty} P_n(x) t^n,$$

where $A(t)$ is a formal power series in t with $A(0) = 1$. This definition implies the equivalent property that

$$(1.2) \quad DP_n(x) = P_{n-1}(x), \quad D = d/dx,$$

Examples of such polynomial sets are

$$(1.3) \quad \left\{ \frac{x^n}{n!} \right\}, \left\{ \frac{B_n(x)}{n!} \right\}, \left\{ \frac{H_n(x)}{2^n n!} \right\}$$

where $B_n(x)$ is the n th Bernoulli polynomial and $H_n(x)$ is the n th Hermite polynomials generated by

$$(1.4) \quad e^{2xt-t^2} = \sum_{n=0}^{\infty} H_n(x) \frac{t^n}{n!}.$$

By an orthogonal polynomial set (OPS) we shall mean those polynomial sets which satisfy a three term recurrence relation of the form

1991 *Mathematics Subject Classification.* Primary 33D45, 33D05; Secondary 42A65.

Key words and phrases. Orthogonal polynomials, generating functions, Askey-Wilson operator.

Research was partially supported by NSERC (Canada) grant A2975.

$$(1.5) \quad P_{n+1}(x) = (A_n x + B_n)P_n(x) - C_n P_{n-1}(x), \quad (n = 0, 1, 2, \dots)$$

with $P_0(x) = 1$, $P_{-1}(x) = 0$, and $A_n A_{n-1} C_n > 0$.

By Favard's theorem [7] this is equivalent to the existence of a positive measure $d\alpha(x)$ such that

$$(1.6) \quad \int_{-\infty}^{\infty} P_n(x) P_m(x) d\alpha(x) = K_n \delta_{nm}.$$

As we see from the examples (1.3) some Appell polynomials are orthogonal and some are not. This prompted Angelesco [3] to prove that *the only orthogonal polynomial sets which are also Appell is the Hermite polynomial set*. This theorem was rediscovered by several authors later on (see, e.g., [10]).

There were several extensions and/or analogs of Appell polynomials that were introduced later. Some are based on changing the operator D in (1.2) into another differentiation-like operator or by replacing the generating relation (1.1) by a more general one. In most of these cases theorems like Angelesco's were given. For example Carlitz [6] proved that the Charlier polynomials are the only OPS which satisfy the difference relation

$$(1.7) \quad \Delta P_n(x) = P_{n-1}(x), \quad (\Delta f(x) = f(x+1) - f(x).)$$

See [1] for many other references.

A new and very interesting analog of Appell polynomials were introduced recently, as a biproduct of other considerations, by Ismail and Zhang [9]. In discussing the Askey-Wilson operator they defined a new q -analog of the exponential function e^{xt} . This we describe in the next section.

2. NOTATIONS AND DEFINITIONS

The Askey-Wilson operator is defined by

$$(2.1) \quad \mathcal{D}_q f(x) = \frac{\delta_q f(x)}{\delta_q x},$$

where $x = \cos \theta$ and

$$(2.2) \quad \delta_q g(e^{i\theta}) = g(q^{1/2} e^{i\theta}) - g(q^{-1/2} e^{i\theta}).$$

We further assume that $-1 < q < 1$ and use the notation

$$(2.3) \quad (a; q)_0 = 1, \quad (a; q)_n = (1-a)(1-qa) \cdots (1-aq^{n-1}), \quad (n = 1, 2, \dots)$$

$$(2.4) \quad (a; q)_\infty = \prod_{k=0}^{\infty} (1-aq^k).$$

There are two q -analogs of the exponential function e^x given by the infinite products

$$(2.5) \quad e_q(x) = \frac{1}{(x; q)_\infty} = \sum_{k=0}^{\infty} \frac{x^k}{(q; q)_k},$$

and

$$(2.6) \quad \frac{1}{e_q(x)} = (x; q)_\infty = \sum_{k=0}^{\infty} (-1)^k \frac{q^{\frac{1}{2}k(k-1)}}{(q; q)_k} x^k.$$

We shall also use the function

$$(2.7) \quad \Psi_n(x) = i^n (iq^{(1-n)/2} e^{i\theta}; q)_n (iq^{(1-n)/2} e^{-i\theta}; q)_n,$$

so that

$$(2.8) \quad \begin{aligned} \Psi_{2n}(x) &= \prod_{k=0}^{n-1} \left[4x^2 + (1 - q^{2n-1-2k})(1 - q^{1-2n+2k}) \right] \\ \Psi_{2n+1}(x) &= 2x \prod_{k=0}^{n-1} \left[4x^2 - (1 - q^{2n-2k})(1 - q^{-2n+2k}) \right] \\ 4x^2 \Psi_n(x) &= \Psi_{n+2}(x) + (1 - q^{n+1})(1 - q^{-n-1}) \Psi_n(x) \end{aligned}$$

Thus

$$(2.9) \quad \mathcal{D}_q \Psi_n(x) = 2q^{(1-n)/2} \frac{1 - q^n}{1 - q} \Psi_{n-1}(x).$$

and

$$(2.10) \quad \mathcal{D}_q [x \Psi_n(x)] = \frac{q^{(1+n)/2} - q^{-(n+1)/2}}{q^{1/2} - q^{-1/2}} 2x \Psi_{n-1}(x).$$

Iterating (2.9) we get

$$(2.11) \quad \mathcal{D}_q^k \Psi_n(x) = 2^k q^{\frac{1}{4}k(k+1) - \frac{1}{2}nk} \frac{(q; q)_n}{(q; q)_{n-k} (1 - q)^k} \Psi_{n-k}(x).$$

The Ismail-Zhang q -analog of the exponential function [9] is

$$(2.12) \quad \mathcal{E}(x) = \sum_{n=0}^{\infty} \frac{q^{n(n-1)/4} (1 - q)^n}{2^n (q; q)_n} \Psi_n(x) t^n.$$

It follows from (2.12) and (2.9) that

$$(2.13) \quad \mathcal{D}_q \mathcal{E}(x) = t \mathcal{E}(x).$$

This suggested to Ismail and Zhang to define the $\mathcal{D}_q$ -Appell polynomials as those, in analogy with (1.1), defined by

$$(2.14) \quad A(t) \mathcal{E}(x) = \sum_{n=0}^{\infty} P_n(x) t^n,$$

so that

$$(2.15) \quad \mathcal{D}_q P_n(x) = P_{n-1}(x).$$

An example of such a set is the Rogers q -Hermite polynomials, $\{H_n(x|q)\}$, (see [2, 4, 8]).

$$(2.16) \quad \prod_{n=0}^{\infty} (1 - 2xtq^n + t^2 q^{2n})^{-1} = \sum_{n=0}^{\infty} H_n(x|q) \frac{t^n}{(q; q)_n}.$$

They satisfy the three term recurrence relation

$$(2.17) \quad H_{n+1}(x|q) = 2xH_n(x|q) - (1 - q^n)H_{n-1}(x|q), \quad n = 0, 1, 2, 3, \dots$$

with $H_0(x|q) = 1$, $H_{-1}(x|q) = 0$.

3. THE MAIN RESULT

We now state our main result:

Theorem 1. *The orthogonal polynomial sets which are also $\mathcal{D}_q$ -Appell, i.e., satisfy (2.15) or (2.14) is the set of the Rogers q -Hermite polynomials.*

Proof Let $\{Q_n(x)\}$ be a polynomial set which is both orthogonal and $\mathcal{D}_q$ -Appell. That is $\{Q_n(x)\}$ satisfy (2.14) and (1.5).

We next note that (2.16) implies that

$$(3.1) \quad h_n(x|q) = \frac{(1 - q)^n q^{n(n-1)/4}}{2^n (q; q)_n} H_n(x|q)$$

satisfy

$$(3.2) \quad \mathcal{D}_q h_n(x|q) = h_{n-1}(x|q),$$

so that $\{h_n(x|q)\}$ is a $\mathcal{D}_q$ -Appell polynomial set and at the same time is an OPS satisfying the three term recurrence relation

$$(3.3) \quad (1 - q^{n+1})h_{n+1}(x|q) = (1 - q)q^{n/2}xh_n(x|q) - \frac{1}{4}(1 - q)^2q^{n-1/2}h_{n-1}(x|q)$$

It also follows from (2.14) that any two polynomial sets $\{R_n(x)\}$ and $\{S_n(x)\}$, in that class are related by $R_n(x) = \sum_{k=0}^n c_{n-k} S_k(x)$. Thus the solution to our problem may be expressed as

$$(3.4) \quad Q_n(x) = \sum_{k=0}^n a_{n-k} h_k(x|q).$$

for some sequence of real constants $\{a_n\}$. We may assume without loss of generality that $a_0 = 1$.

The three term recurrence relation satisfied by $\{Q_n(x)\}$ is

$$(3.5) \quad (1 - q^{n+1})Q_{n+1}(x) = ((1 - q)q^{n/2}x + \beta_n)Q_n(x) - \gamma_n Q_{n-1}(x),$$

with $Q_0(x) = 1$, $Q_{-1}(x) = 0$. Thus $Q_1(x) = x + \beta_0 = a_1 + h_1(x|q)$, from which it follows that $a_1 = \beta_0$.

Putting (3.4) in (3.5) and using (3.3) to replace $xh_k(x|q)$ in terms of $h_{k+1}(x|q)$ and $h_{k-1}(x|q)$ we get, on equating coefficients of $h_k(x|q)$,

$$(3.6) \quad (1 - q^{(n-k+1)/2})(1 + q^{(n+1+k)/2})a_{n+1-k} - \beta_n a_{n-k} + \left[\gamma_n - \frac{1}{4}(1 - q)^2 q^{(n+k)/2} \right] a_{n-k-1} = 0,$$

valid for all n and $k = 0, 1, 2, \dots, n+1$ provided we interpret $a_{-1} = a_{-2} = 0$. It is easy to see that this system of equations is equivalent to the solution of our problem.

Putting $k = n$ in (3.6) we get

$$(3.7) \quad \beta_n = (1 - q^{\frac{1}{2}})(1 + q^{n+\frac{1}{2}})a_1.$$

Hence if $\beta_0 = 0$ then $\beta_n = 0$ for all n . In fact if $\beta_m = 0$ for any $n = m$ then $\beta_n = 0$ for all n .

Now we treat these two cases separately.

Case I. ($\beta_0 = 0$).

The system (3.6) can now be written as

$$(3.8) \quad (1 - q^{(k+1)/2})(1 + q^{n+\frac{1}{2}(1-k)})a_{k+1} + \left[\gamma_n - \frac{1}{4}(1 - q)^2 q^{n-\frac{1}{2}k} \right] a_{k-1} = 0.$$

Since $a_1 = 0$ then it follows from (3.8) that $a_{2k+1} = 0$ for all k . In particular we get

$$(3.9) \quad \gamma_n = \frac{1}{4}(1 - q)^2 q^{n-\frac{1}{2}} - a_2(1 - q)(1 + q^n),$$

so that if $a_2 = 0$ then

$$(3.10) \quad Q_n(x) = h_n(x|q).$$

Now we show that $a_2 \neq 0$ leads to contradiction. To do this replace k by $2k - 1$. We get

$$(3.11) \quad (1 - q^k)(1 + q^{n-k+1})a_{2k} + \left[\frac{1}{4}(1 - q)^2 q^{n-\frac{1}{2}}(1 - q^{1-k}) - a_2(1 - q)(1 + q^n) \right] a_{2k-2} = 0.$$

Keep k fixed and let $n \rightarrow \infty$. We get $(1 - q^k)a_{2k} = (1 - q)a_2a_{2k-2}$. Thus

$$(3.12) \quad a_{2k} = \frac{(1 - q)^k}{(q; q)_k} a_2^k.$$

Putting this value in (3.11) we get $q^{1-k} = 1$. This is a contradiction and Case I is finished.

Case II ($\beta_0 \neq 0$).

We start with (3.6) we get, assuming $a_1 \neq 0$,

$$(3.13) \quad \gamma_n = \frac{1}{4}(1 - q)^2 q^{n-\frac{1}{2}} + (1 - q^{\frac{1}{2}})(1 + q^{n+\frac{1}{2}})a_1^2 - (1 - q)(1 + q^n)a_2.$$

Putting this value of γ_n and the value of β_n in (3.7) in (3.6), and finally equating coefficients of q^n and the terms independent of n we get the pair of equation systems

$$(3.14) \quad (1 - q^{(k+1)/2})a_{k+1} - (1 - q^{\frac{1}{2}})a_1a_k + \left\{ (1 - q^{\frac{1}{2}})a_1^2 - (1 - q)a_2 \right\} a_{k-1} = 0$$

and

$$(3.15) \quad (1 - q^{(k+1)/2})a_{k+1} - (1 - q^{\frac{1}{2}})q^{k/2}a_1a_k + \left\{ \frac{1}{4}(1 - q)^2 q^{-\frac{1}{2}}(q^{(k-1)/2} - 1) + q^{k/2}(1 - q^{\frac{1}{2}})a_1^2 - (1 - q)q^{(k-1)/2}a_2 \right\} a_{k-1} = 0$$

Eliminating a_{k+1} in these equations we get

$$(3.16) \quad (1 - q^{\frac{1}{2}})(1 - q^{k/2})a_1a_k + \left\{ (1 - q)a_2(1 - q^{(k-1)/2}) - (1 - q^{\frac{1}{2}})(1 - q^{k/2})a_1^2 - \frac{1}{4}(1 - q)^2 q^{-\frac{1}{2}}(1 - q^{(k-1)/2}) \right\} a_{k-1} = 0.$$

This equation is of the form $(1 - q^{k/2})a_1a_k = c(1 - bq^{k/2})a_{k-1}$ so that the general solution of (3.16) is

$$(3.17) \quad a_k = c^k \frac{(bq^{\frac{1}{2}}; q^{\frac{1}{2}})_k}{(q^{\frac{1}{2}}; q^{\frac{1}{2}})_k}$$

Putting this in (3.14) we get that $b = 0$. On the other hand (3.15) gives that $c^2 = \frac{1}{4}(1 - q)^2 q^{-\frac{1}{2}}$. Finally putting those values of a_k in (3.13) we get that $\gamma_n = 0$ which is a contradiction.

This completes the proof of the theorem.

4. GENERATING FUNCTION

We obtain, for the q-Hermite polynomials, a generating function of the form (2.14). More specifically we prove

Theorem 2. *Let $H_n(x|q)$ be the n th Rogers q -Hermite polynomial. Then we have*

$$(4.1) \quad \sum_{n=0}^{\infty} \frac{q^{n(n-1)/4}}{(q; q)_n} H_n(x|q) t^n = (t^2 q^{-\frac{1}{2}}; q^2)_{\infty} \mathcal{E}(x).$$

Proof. Let $A(t) = 1 + a_1 t + a_2 t^2 + a_3 t^3 + \dots$ and

$$(4.2) \quad A(t) \mathcal{E}(x) = \sum_{n=0}^{\infty} h_n(x|q) t^n.$$

Then we get

$$(4.3) \quad h_n(x|q) = \sum_{k=0}^n a_{n-k} c_k \Psi_k(x).$$

where

$$(4.4) \quad c_k = \frac{(1-q)^k}{2^k (q; q)_k} q^{k(k-1)/4}.$$

To calculate the coefficients $\{a_n\}$ we first iterate (3.3) we get

$$(4.5) \quad \begin{aligned} 4x^2 h_n(x|q) &= \frac{4}{(1-q)^2} (1-q^{n+1})(1-q^{n+2}) q^{-n-\frac{1}{2}} h_{n+2}(x|q) \\ &+ (2-q^n-q^{n+1}) h_n(x|q) + \frac{(1-q)^2}{4} q^{n-\frac{3}{2}} h_{n-2}(x|q). \end{aligned}$$

Putting (4.3) in (4.5), using (2.6) and then equating coefficients of $\Psi_k(x)$ we get after some simplification

$$(4.6) \quad \begin{aligned} &\frac{4}{(1-q)^2} q^{-n-\frac{1}{2}} (1-q^{n-k+2})(1-q^{n+k+1}) a_{n+2-k} + \\ &q^{-k-1} \{1 + q^{2k+2} - q^{n+k+1} - q^{n+k+2}\} a_{n-k} + \\ &\frac{(1-q)^2}{4} q^{n-\frac{3}{2}} a_{n-2-k} = 0 \quad (k = 0, 1, \dots, n+2). \end{aligned}$$

By direct calculation of a_1, a_2, a_3 we see easily that $a_1 = a_3 = 0$. Thus (4.6) shows that $a_{2k+1} = 0$ for all k .

Furthermore we can easily verify that

$$(4.7) \quad a_{2j} = (-1)^j \frac{(1-q)^{2j}}{2^{2j} (q^2; q^2)_j} q^{j(j-\frac{3}{2})} \quad (j = 0, 1, 2, 3, \dots)$$

Hence

$$\begin{aligned}
(4.8) \quad A(t) &= \sum_{j=0}^{\infty} (-1)^j \frac{q^{j(j-1)}}{(q^2; q^2)_j} \left(\frac{(1-q)^2 t^2}{4} q^{-\frac{1}{2}} \right)^j \\
&= \left(\frac{(1-q)^2}{4} t^2 q^{-\frac{1}{2}}; q^2 \right)_{\infty}.
\end{aligned}$$

After some rescaling we get the theorem.

As a corollary of (4.1) we state the pair of inverse relations

$$(4.9) \quad \Psi_n(x) = \sum_k \frac{(q; q)_n q^{k(k-n)}}{(q^2; q^2)_k (q; q)_{n-2k}} H_{n-2k}(x|q),$$

$$(4.10) \quad H_n(x|q) = \sum_k (-1)^k \frac{(q; q)_n q^{k(2k-n-1)}}{(q^2; q^2)_k (q; q)_{n-2k}} \Psi_{n-2k}(x).$$

These follow from the identities (2.5) and (2.6)

Formula (4.10) and (2.11) give

$$(4.11) \quad H_n(x|q) = \frac{1}{e_{q^2}\left(\frac{(1-q)^2}{4} q^{-\frac{1}{2}} \mathcal{D}_q^2\right)} \Psi_n(x).$$

This is a q-analog of the formula

$$e^{-D^2} x^n = H_n(x)$$

for the regular Hermite polynomials (1.4).

REFERENCES

- [1] W. A. Al-Salam, *Characterization theorems for orthogonal polynomials*, in "Orthogonal Polynomials: Theory and Practice", P. Nevai ed., Kluwer, Dordrecht, 1989, pp. 1–24.
- [2] W. A. Al-Salam and M. E. H. Ismail, q-beta integrals and the q-Hermite polynomials, *Pacific J. of Math.*, **135**(1988), 209–221.
- [3] A. Angelesco, Sur les polynomes orthogonaux en rapport avec d'autre polynomes, *Buletinul Societății din Cluj*, **1**(1921), 44–59.
- [4] R. Askey and M. E. H. Ismail, *A generalization of the ultraspherical polynomials*, *Studies in Pure Mathematics*, edited by P. Erdős, pp. 55–78, Birkhäuser, Basel, 1983.
- [5] R. Askey and J. Wilson, *Some basic hypergeometric polynomials that generalize Jacobi polynomials*, *Memoirs Amer. Math. Soc.* #319, Providence, 1985.
- [6] L. Carlitz, *Characterization of certain sequences of orthogonal polynomials*, *Portugaliae Math*, **20**(1961), 43–46.
- [7] T. S. Chihara, *An Introduction to Orthogonal Polynomials*, Gordon and Breach, 1978.
- [8] M. E. H. Ismail and D. Stanton, *On the Askey-Wilson and Rogers polynomials*, *Canad. J. Math.* **40** (1988), 1025–1045.
- [9] M. E. H. Ismail and R. Zhang, *Diagonalization of certain integral operators*, *Advances in Mathematics*, to appear.

- [10] J. Shohat, *The relation of the classical orthogonal polynomials to the polynomials of Appell*, American J. Math., **58**(1936), 453-464.

(Waleed. A. Al-Salam) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ALBERTA, EDMONTON,
CANADA, T6G 2G1

E-mail address: waleed@euler.math.ualberta.ca